Card No BAT03-IT

Review of the 17/05/2024

Aleabacter HA

COMPOSITION

Pre-acclimatized and freeze-dried culture of Oenococcus oeni

GENERAL CHARACTERISTICS

aLEABACTER HA is a culture of *Oenococcus oeni* bacteria selected for proper management of malolactic fermentation (MLF) in musts and wines. The specific bacterial strain was selected for wines with high fixed acidity and low pH.

During production, the bacterial culture is acclimated to very low pH conditions and is therefore already prepared for these difficult conditions.

Parameters of use (NB: attention the parameters act synergistically):

ALCOL	TEMPERATURA	SO ₂	Ph
> 14,5 % vol.	17-23 °C	< 8 mg/L SO ₂ libera (< ~30 mg/L SO ₂ totale)	> 3.00

TECHNICAL SPECIFICATIONS

Total bacteria counts: $> 1 \times 10^{1011}$ U.F.C. / g

Molds: $< 1 \times 10^{103}$ U.F.C. / g

Yeasts: $< 1 \times 10^{3} \text{ U.F.C.} / \text{ g}$

Acetobacteria: $< 1 \times 10^{4}$ U.F.C. / g

E. coli: absent

Salmonella: absent in 10 g

Listeria: absent in 25 g

APPLICATIONS

ALEABACTER HA is recommended for initiating MLF in both white and red wines that have low pH values (< 3.30). It can be used:

- with classic sequential inoculation, at the end of alcoholic fermentation, after an initial racking to remove coarse lees;
- in co-inoculum after 24-48 h after inoculation of selected yeasts on must.

In co-inoculation it was tested with positive results with the following yeasts from the Aleaferm range: Reserve, 0-12.

MODE OF USE

Rehydrate **ALEABACTER HA** in mineral water (DO NOT use tap water, demineralized or distilled water) at a ratio of 1:20 (0.5 L for a 25 hL bag; 5 L for a 250 hL bag), at about 25° C for 10-15 minutes while stirring gently.

Then add them to the mass to be treated according to the following directions:

CO-INOCULUM at 24-48 h after Saccharomyces yeast inoculum:

- It is recommended to add to a fraction of the must to be inoculated and allow to acclimatize for 2-3 hours. Add the suspension directly to the mass to be treated. In case of maceration make the addition under cap.
- No pumping over or agitation is necessary as normal fermentation activity is sufficient to homogenize the bacterial culture.
- It is recommended to keep the fermentation temperature below 28°C. If the temperature exceeds 30°C there will be a halt in malic acid consumption, which will resume independently as the temperature falls into the optimal ranges.



40062 Molinella (BO) - ITALIA
Tel. +39 051 88 7052
+39 346 58 33 882
info@alea-evolution.com
alea-evolution@pec.it
alea-evolution.com
CF - P.IVA 02944251202
Cod. REA BO-479327

Via Sandro Pertini, 12



Capitale sociale € 40.000,00 i.v.



The information given in this Technical Data Sheet corresponds to the current state of our knowledge and is subject to modification and supplementation without prior notice. The methods of use given do not relieve the user from the application and observance of safety and protection regulations. Adaptation to individual cases, as a consequence of the specific circumstances of each use, as well as possible misuse of the product, do not involve the responsibility of Alea Evolution S.R.L.

Card No BAT03-IT

Review of the 17/05/2024

Analytical monitoring, preferably daily, is recommended to verify the proper progress of fermentations. **EQUENTIAL inoculation**at the end of AF:

At least one racking is recommended to remove coarse lees and inoculate on clean wine to further limit the formation of volatile acidity and reduce the risk of olfactory deviations.

We recommend increasing the acclimatization time before inoculation to 6-12 hours (in any case, do not exceed 24 hours) taking care to use a sufficiently high volume of wine to allow the bacterial culture not to go into competitive stress given the high cell concentration.

The use of the specific activator **leavit FML**is recommended in any case, but is essential in case of very clear wines and in general in case of late inoculation (e.g., the spring following AF).

Do not make $_{\rm O2}$ additions at the end of AF, before inoculation of bacteria and during the course of MLF. Simultaneous use of β -glucanase does not affect the smooth conduct of MLF.

nalytical control of MLF with analysis every 2-3 days is recommended. As soon as the residual value of ac. L-malic acid falls below 0.2-0.3 g/L, addition of $_{O2}$ and/or cooling of the mass to 8-10 $^{\circ}$ C is recommended to avoid consumption of citric acid with possible production of acetic acid and diacetyl.

PACKAGES AVAILABLE

STORAGE CONDITIONS

Store in unopened package in the freezer at a temperature below -18° C (for short periods before use, storage at $+4^{\circ}$ C is possible). Thawed product must be used quickly and CANNOT be refrozen.

Exclusively for oenological and professional use - Reg. (EU) 2022/68

Product obtained from raw materials in accordance with O.I.V. International Oenological Codex



Tel. +39 051 88 7052 +39 346 58 33 882 info@alea-evolution.com alea-evolution@pec.it alea-evolution.com CF - P.IVA 02944251202 Cod. REA BO-479327 Capitale sociale € 40.000,00 i.v.

Via Sandro Pertini, 12 40062 Molinella (BO) - ITALIA





The information given in this Technical Data Sheet corresponds to the current state of our knowledge and is subject to modification and supplementation without prior notice. The methods of use given do not relieve the user from the application and observance of safety and protection regulations. Adaptation to individual cases, as a consequence of the specific circumstances of each use, as well as possible misuse of the product, do not involve the responsibility of Alea Evolution S.R.L.